$$\Delta y = \left[\frac{v_{y_0} + v_y}{2}\right]t$$

$$\Delta y = v_{y_0} \Delta t + \frac{1}{2} a_y t^2$$

$$v_y = v_{y_0} + a_y t$$

$$v_y^2 = v_{y_0}^2 + 2a_y \Delta y$$

$$g = 9.8m/s^2$$

$$ax^2 + bx + c = 0$$

$$\mathbf{x} = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$m \rightarrow 10^{-3}$$

$$k \rightarrow 10^3$$

$$1m = 3.28 ft$$

$$a_{x} = 0$$

$$\Delta x = \left[\frac{v_{x_0} + v_x}{2}\right] t$$

$$\Delta x = v_{x_0} t + \frac{1}{2} a_x t^2$$

$$\Delta x = v_{x_0} t + \frac{1}{2} a_x t^2$$

$$v_{x} = v_{x_0} + a_{x}t$$

$$v_x^2 = v_{x_0}^2 + 2a_x \Delta x$$

$$|\vec{A}| = \sqrt{A_x^2 + A_y^2}$$

$$\sin\theta = \frac{A_y}{|\vec{A}|}$$

$$\cos\theta = \frac{A_x}{|\vec{A}|}$$

$$\tan \theta = \frac{A_y}{A_x}$$



Show all work in the spaces provided.


1) Let

$$|\vec{A}| = 3.0 \, m \, \text{at} \, \theta_A = 160^{\circ}$$
  
 $|\vec{B}| = 2.0 \, m \, \text{at} \, \theta_B = 0^{\circ}$ 

$$\left| \vec{B} \right| = 2.0 \, m \, \text{at} \, \theta_{\scriptscriptstyle B} = 0^{\circ}$$

$$\left| \vec{C} \right| = 5.0 \, m \, \text{at} \, \theta_C = 210^\circ$$

a) Draw the vectors on a coordinate system using the graph paper below. (5 points)



b) What are the x and y components of Vectors 
$$\vec{A}, \vec{B}$$
 and  $\vec{C}$ ? (6 points)

 $A_x = |\vec{A}| Coo (160^\circ)$ 
 $A_x = 3 N Coo (160^\circ)$ 
 $A_x = -2.82 M$ 
 $A_x = -2.82 M$ 
 $A_x = -3.82 M$ 
 $A_x$ 

$$Ay = 1.03m$$

$$\beta_{x} = 2m$$

$$\beta_{y} = 0$$

$$C_{x} = -4.33 \text{ m}$$

$$c_g = -2.5 \text{ m}$$

c) What is the sum of Vectors  $\vec{A}, \vec{B}$  and  $\vec{C}$  in terms of its magnitude and direction? (4 points)

$$R_x = A_x + B_x + C_x$$

$$R_x = -2.82m + 2m - 4.33m$$

$$R_x = -5.15 \,\mathrm{m}$$

$$\hat{\beta} = (-5.15 \text{ m}, -1.47 \text{ m})$$

$$|\vec{R}| = \sqrt{R_x^2 + R_y^2}$$

$$|\vec{R}| = (-5.15 \,\mathrm{m})^2 + (-1.47 \,\mathrm{m})^2$$

$$|R| = 5.36 \,\mathrm{m}$$


$$\Theta = tan^{-1} \left( \frac{R_y}{R_x} \right)$$

Ry = Ay + By + Cy

 $\hat{R}_{x} = -1.47 \text{ m}$ 

 $R_y = 1.03m + 0 - 2.5m$ 

Test 1 Review Page 5



- 2) A cannonball aimed at  $40^{\circ}$  is fired at a wall  $300\,m$  away on level ground as shown above. The initial speed of the cannonball is  $|\vec{v}_o| = 89 \, m/s$  .

a) How long does it take the cannonball to hit the wall? (3 points)
$$\Delta X = V_{x} t$$

$$t = \frac{\Delta X}{V_{x}} = \frac{300 \text{M}}{(3.13 \text{ m/s})} = 4.45$$

b) At what height does it hit the wall? (4 points)
$$\Delta g = V_1 \times + \pm \Delta g \times$$

$$\Delta g = (57.2 | m|_8)(4.48) + \pm (-9.8 | m|_8^2)(4.48)^2$$

$$\Delta g = 251.72 | m - 94.36 | m = 156.86 | m$$
c) What is the speed and direction when it hits the wall? Is it on the way up or down? (4 points)
$$V_1 = 6.9.19 | m|_8$$

Dx = 68, 18 m/s

$$V_y = V_y + a_y t$$
  
 $V_y = 57.21 \text{ m/s} + (-9.8 \text{ m/s}^2)(4.45)$   
 $V_y = 14.09 \text{ m/s}$ 

$$0 = 0$$
 $0 = 9.8 \text{m/s}^2$ 
 $0 = 9.8 \text{m/s}^2$ 
 $0 = 10.1 \text{ Cas} (40^\circ)$ 
 $0 = (89 \text{m/s}) \text{ Cas} (40^\circ)$ 
 $0 = (89 \text{m/s}) \text{ Sin} (40^\circ)$ 

$$|\vec{y}| = |y_x^2 + y_y^2| |\vec{y}| = |(68.18 \text{ m/s})^2 + (14.09 \text{ m/s})^2| |\vec{y}| = 69.62 \text{ m/s} (Spece) \Theta = tun^2 (\frac{y_y}{y_x}) \Theta = tun^4 (\frac{14.09 \text{ m/s}}{68.18 \text{ m/s}}) = 11.680 (dig.)$$

